



DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

MOUSEOFF® ZP ECONOBAIT®

FOR EMERGENCY USE ONLY

ACTIVE CONSTITUENT: 25 g/kg Zinc Phosphide

For the reduction of heavy infestations of mice in agricultural situations.

THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROLS FOR SCHEDULE 7 POISONS.

Net contents: 125 kg

DIRECTIONS FOR USE:

RESTRAINTS:

DO NOT apply to bare ground (including fallow where there is no vegetative cover).

DO NOT apply in a trail.

DO NOT apply if heavy rain is imminent

DO NOT apply to the outer 50m of crop or within 50m of native vegetation. A 50m buffer zone in crop must be employed to reduce risk to birds and other non-target animals.

DO NOT apply unless monitoring of mouse numbers indicates the potential for crop damage of economic significance.

PROTECTION OF WILDLIFE, THE ENVIRONMENT AND LIVESTOCK:

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath the aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

STORAGE AND DISPOSAL:

The contents of each container should be used within 3 months of opening. Store in the closed original container in a dry cool well-ventilated area out of direct sunlight. Store in a locked area away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition. Dispose of used containers by crushing and burying below one meter in a disposal pit specifically marked and set up for this purpose, clear of waterways, vegetation and roots. Excess or unused baits must be buried below one metre. Empty containers and product must not be burnt.

SAFETY DIRECTIONS:

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames – forms toxic gas. When opening the container and using baits wear elbow-length PVC gloves and a full face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each day's use wash gloves and respirator and if rubber wash with detergent and warm water.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. Do not give direct mouth-to-mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy-viva or one-way mask. Resuscitate in a well ventilated area.

MSDS:

Additional information is listed in the MSDS which can be obtained from the supplier. www.animalcontrol.com.au

CONDITIONS OF SALE and DISCLAIMER

Animal Control Technologies Pty Ltd (ACTA) will not accept responsibility for losses or damage arising from the supply or use of these goods other than the responsibility for the merchantable quality of the goods. To the extent allowed by the relevant laws in each State, the liability of ACTA is limited to the replacement of goods in the event that a valid claim of deficiencies in merchantable quality is proven.

SITUATION	PEST	RATE	CRITICAL COMMENTS
Grain crops, legume crops, canola, safflower, nut crops, and pasture. (Note: for all situations – only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seed has been sown into crop stubble, into cultivated ground or other similar ground cover).	Mice	1 kg/ha to achieve an even coverage of 2-3 grains/m ²	Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2-3 grains/m ² . It is imperative all attempts are made to apply bait as evenly as possible to maximize product efficacy and minimize risks to non-target animals. MOUSEOFF® ZP ECONOBAIT® will provide a high level of control in most situations. However in areas of extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Retreatment may therefore be required in certain circumstances. It is recommended that pre-treatment estimations of the mouse problem severity is performed before applying bait and post-baiting estimates are conducted to assess the effectiveness of the control operation. Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION. THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN

APVMA Permit No: PER12899

BATCH:

DOM: